

Dear Parent

The UK Health Security agency are reporting that we are seeing a higher number of cases of group A strep this year than usual in the North West. The bacteria usually causes a mild infection producing sore throats or scarlet fever that can be easily treated with antibiotics.

In very rare circumstances, this bacteria can get into the bloodstream and cause serious illness – called invasive group A strep (iGAS). This is still uncommon however it is important that parents are on the lookout for symptoms and see a doctor as quickly as possible so that their child can be treated and we can stop the infection becoming serious.

Make sure you talk to a health professional if your child is showing signs of deteriorating after a bout of scarlet fever, a sore throat, or a respiratory infection.

Scarlet fever is usually a mild illness, but it is highly infectious. Therefore, look out for symptoms in your child, which include a sore throat, headache, and fever, along with a fine, pinkish or red body rash with a sandpapery feel. On darker skin the rash can be more difficult to detect visually but will have a sandpapery feel.

Contact NHS 111 or your GP if you suspect your child has scarlet fever, because early treatment of scarlet fever with antibiotics is important to reduce the risk of complications such as pneumonia or a bloodstream infection. If your child has scarlet fever, keep them at home until at least 24 hours after the start of antibiotic treatment to avoid spreading the infection to others.

There are lots of viruses that cause sore throats, colds and coughs circulating. These should resolve without medical intervention. However, children can on occasion develop a bacterial infection on top of a virus and that can make them more unwell.

As a parent, if you feel that your child seems seriously unwell, you should trust your own judgement. Contact NHS 111 if or your GP if:

- your child is getting worse
- your child is feeding or eating much less than normal
- your child has had a dry nappy for 12 hours or more or shows other signs of dehydration
- your baby is under 3 months and has a temperature of 38C, or is older than 3 months and has a temperature of 39C or higher
- your baby feels hotter than usual when you touch their back or chest, or feels sweaty
- your child is very tired or irritable

Call 999 or go to A&E if:

- your child is having difficulty breathing – you may notice grunting noises or their tummy sucking under their ribs
- there are pauses when your child breathes
- your child's skin, tongue or lips are blue
- your child is floppy and will not wake up or stay awake

Good hand and respiratory hygiene are important for stopping the spread of many bugs. By teaching your child how to wash their hands properly with soap for 20 seconds, using a tissue to catch coughs and sneezes, and keeping away from others when feeling unwell, they will be able to reduce the risk of picking up, or spreading, infections.

More information can be found here: [Scarlet fever: symptoms, diagnosis and treatment - GOV.UK \(www.gov.uk\)](https://www.gov.uk/government/healthcare-articles/scarlet-fever-symptoms-diagnosis-and-treatment)